arXiv:1011.0535v2 [math.CV] 4 Nov 2010

CARATHEODORY CONVERGENCE OF LOG-RIEMANN SURFACES AND EULER'S FORMULA

KINGSHOOK BISWAS AND RICARDO PEREZ-MARCO

Abstract. We define the notion of log-Riemann surfaces and Caratheodory convergence of log-Riemann surfaces. We prove a convergence theorem for uniformizations of simply connected log-Riemann surfaces converging in the Caratheodory topology. We obtain as a corollary a purely geometric proof of Euler's formula $\left(1+\frac{z}{n}\right)^n \to e^z$.

Contents

1. Introduction 1
1.1. Acknowledgements 2
2. Log-Riemann surfaces 3
2.1. Definitions 3
2.2. Examples 4
3. Caratheodory Theorem for log-Riemann surfaces 5
3.1. Caratheodory convergence of log-Riemann surfaces 5
3.2. Convergence of uniformizations 6
References 7

1. Introduction

In this article we introduce the notion of *log-Riemann surfaces*. These are Riemann surfaces informally described as obtained by cutting and pasting copies of $\mathbb{C}$ isometrically. The larger class of *tube-log Riemann surfaces*, allowing pasting of flat cylinders $\mathbb{C}/\mathbb{Z}$ as well, was used by the second author for the construction of holomorphic diffeomorphisms with irrationally indifferent fixed points with special dynamical properties, see [PM93], [PM95], [PM00]. They were also used by the first author to construct further examples in [Bis05], [Bis08]. Our aim in this article is to initiate a general study of log-Riemann surfaces.

A log-Riemann surface $\mathcal{S}$ comes equipped with a local diffeomorphism $\pi : \mathcal{S} \to \mathbb{C}$ satisfying certain conditions (defined precisely in the next section). The map π serves as a coordinate at all points of $\mathcal{S}$ and allows one to write formulae for the uniformizations of simply connected log-Riemann surfaces. The map π also induces a flat metric on $\mathcal{S}$; the points p added in the completion are "ramification points" of π, i.e. π restricted to a small punctured disc neighbourhood of p is a covering of a punctured disc in $\mathbb{C}$. The order of a ramification point is defined to be the degree $1 \leq n \leq \infty$ of the covering; the finite order ramification points can be added to $\mathcal{S}$ to give a Riemann surface $\mathcal{S}^\times$.

For a pointed log-Riemann surface $(\mathcal{S}^{\times}, z_0)$ such that $\mathcal{S}^{\times}$ is simply connected, there is a unique $0 < R \leq +\infty$, called the *conformal radius* of $(\mathcal{S}, z_0)$, and a unique normalized uniformization $F = F_{(\mathcal{S},z_0)} : \mathcal{S}^{\times} \to \mathbb{D}_R$ such that $F(z_0) = 0, F'(z_0) = 1$ (with derivative computed in the chart π). The flat metric also allows one to define a version of Gromov-Hausdorff convergence of pointed log-Riemann surfaces, which we call Caratheodory convergence by analogy with convergence of simply connected domains in $\mathbb{C}$. We then prove an analogue of Caratheodory Kernel Convergence Theorem for log-Riemann surfaces:

Theorem 1.1. *Let* $(\mathcal{S}_n, z_n) \to (\mathcal{S}, z_0)$ *be a Caratheodory convergent sequence of log-Riemann surfaces such that the finite completions* $\mathcal{S}_n^{\times}, \mathcal{S}^{\times}$ *are simply connected. Let the conformal radii and normalized uniformizations of* $(\mathcal{S}_n, z_n), (\mathcal{S}, z_0)$ *be* R_n, R *and* F_n, F *respectively. If* $\limsup_{n\to+\infty} R_n \leq R$, *then* $\lim_{n\to+\infty} R_n = R$ *and the uniformizations* F_n *converge uniformly on compacts of* $\mathcal{S}$ *to the uniformization* F.

The convergence of F_n to F above holds in the following sense: every compact $K \subset \mathcal{S}$ containing z_0 embeds isometrically into $\mathcal{S}_n$ for n large enough via an isometry ι_n which is a translation in the charts π, π_n, and the maps $F_n \circ \iota_n$ converge uniformly to F.

We also have convergence of the inverse mappings assuming the projections of the basepoints converge:

Theorem 1.2. *Suppose* $(\mathcal{S}_n, z_n)$ *converge to* $(\mathcal{S}, z_0)$. *Let* R_n, R *be the corresponding conformal radii and* $G_n : \mathbb{D}_{R_n} \to \mathcal{S}_n^{\times}, G : \mathbb{D}_R \to \mathcal{S}^{\times}$ *be normalized biholomorphisms such that* $G_n(0) = z_n, G(0) = z_0$ *and* $(\pi_n \circ G_n)'(0) = 1, (\pi \circ G)'(0) = 1$. *If* $\limsup R_n \leq R$ *and* $\pi_n(z_n) \to \pi(z_0)$ *then* $\lim R_n = R$ *and* $\pi_n \circ G_n \to \pi \circ G$ *uniformly on compacts of* $\mathbb{D}_R$.

Observing that for the log-Riemann surfaces $\mathcal{S}_n, \mathcal{S}$ corresponding to the nth root and logarithm functions respectively, with base point 1, the normalized uniformizations are given in charts by the functions $\pi_n \circ G_n(z) = (1 + \frac{z}{n})^n$ and $\pi \circ G(z) = e^z$ respectively, we obtain as a direct corollary of the above a purely geometric proof of Euler's formula:

Corollary 1.3. *We have*

$$\left(1 + \frac{z}{n}\right)^n \to e^z$$

uniformly on compacts.

In a forthcoming article [BPM10] we use the above convergence theorems to obtain uniformization formulae for simply connected log-Riemann surfaces with finitely many ramification points.

1.1. **Acknowledgements.** The authors are very grateful to J.Muñoz Díaz, R. Alonso Blanco, J. Lombardero, S. Jim'enez Verdugo, A. Álvarez Vázquez, D. Barsky and F. Marcellan Español for numerous helpful discussions. The first author was partly supported by Department of Science and Technology research project grant DyNo. 100/IFD/8347/2008-2009, and the second author by CNRS, UMR 7539.

2. Log-Riemann surfaces

2.1. **Definitions.**

Definition 2.1. *A log-Riemann surface is a Riemann surface* $\mathcal{S}$ *equipped with a local diffeomorphism* $\pi : \mathcal{S} \to \mathbb{C}$, *called the projection mapping, such that the following holds: let* $\overline{\mathcal{S}} = \mathcal{S} \coprod \mathcal{R}$ *be the completion of* $\mathcal{S}$ *with respect to the path-metric on* $\mathcal{S}$ *induced by the flat metric* $|d\pi|$ *(the pull-back of the Euclidean metric under* π*). Then* $\mathcal{R}$ *is discrete.*

The set $\mathcal{R}$ *is called the ramification locus of* $\mathcal{S}$ *and the points of* $\mathcal{R}$ *are called ramification points.*

Since π is a local isometry with respect to the metrics on $\mathcal{S}$ and $\mathbb{C}$ it extends uniquely to $\overline{\mathcal{S}}$.

Proposition 2.2. *Let* $p \in \mathcal{R}$. *For* $r > 0$ *small enough,* $\pi(B(p,r) - \{p\}) = B(\pi(p), r) - \{\pi(p)\}$ *and*

$$\pi : B(p,r) - \{p\} \to B(\pi(p), r) - \{\pi(p)\}$$

is a covering.

Proof: Choose $r > 0$ small enough so that there are no points of $\mathcal{R}$ in $B(p,r)$ apart from p. Pick a $z \in B(p,r) - \{p\}$; then $\pi(z) \in B(\pi(p), r)$, and the local inverse of π at $\pi(z)$ such that $\pi^{-1}(\pi(z)) = z$ can be analytically continued to all points of $B(\pi(p), r) - \{\pi(p)\}$, since the only possible obstruction to the continuation is encountering points in $\mathcal{R} - \{p\}$ above, but by the choice of r this is not possible. Therefore π maps $B(p,r) - \{p\}$ onto $B(\pi(p), r) - \{\pi(p)\}$. Moreover the local inverse is single-valued on $B(\pi(z), t)$ where $t = d(z,p)$, so π maps $B(z,t)$ isometrically to $B(\pi(z), t)$, so taking $z_n \in B(z,t)$ converging to p, we have

$$|\pi(z) - \pi(p)| = \lim |\pi(z) - \pi(z_n)| = \lim d(z, z_n) = d(z,p) > 0$$

so $\pi(z) \neq \pi(p)$.

The proof that $\pi : B(p,r) - \{p\} \to \pi(B(p,r) - \{p\})$ is a covering is similar. For each point $z_0 \in \pi(B(p,r) - \{p\}) = B(\pi(p), r) - \{\pi(p)\}$, take a small disc $B(z_0, \rho) \subset B(\pi(p), r) - \{\pi(p)\}$ and let U be a connected component of $\pi^{-1}(B(z_0, \rho))$; we can pick a $z_1 \in U$ and as before continue without obstruction the local inverse of π at $\pi(z_1)$ satisfying $\pi^{-1}(\pi(z_1)) = z_1$ to all of the disk $B(z_0, \rho)$; the continuation of π^{-1} to it is single-valued, so $\pi_{|U} : U \to B(z_0, \rho)$ is a diffeomorphism. $\diamond$

Definition 2.3. *The order* $1 \leq n \leq \infty$ *of a ramification point* $p \in \mathcal{R}$ *is defined to be the degree of the covering* $\pi : B(p,r) - \{p\} \to B(\pi(p), r) - \{\pi(p)\}$ *of the punctured disk* $B(\pi(p), r) - \{\pi(p)\}$ *(where* r *is taken small enough as above). The ramification point* p *is called finite if* $n < \infty$ *and infinite if* $n = \infty$.

Definition 2.4. *The finite completion* $\mathcal{S}^{\times} \subset \overline{\mathcal{S}}$ *of a log-Riemann surface* $\mathcal{S}$ *is defined to be the union of* $\mathcal{S}$ *and all finite ramification points of* $\mathcal{S}$.

Since any finite sheeted holomorphic covering of a punctured disc is equivalent to the covering $\mathbb{D}^* \to \mathbb{D}^*, z \mapsto z^n$, a punctured neighbourhood of a finite ramification point is biholomorphic to a punctured disk, so the finite ramification points may be added to $\mathcal{S}$ to give a Riemann surface structure on $\mathcal{S}^\times$ compatible with that of $\mathcal{S}$, such that the map $\pi : \mathcal{S}^\times \to \mathbb{C}$ is holomorphic with critical points at the finite ramification points.

When the finite completion is simply connected, $\mathcal{S}^\times$ is biholomorphic to $\mathbb{D}_R = \{|z| < R\}$ for some $0 < R \leq \infty$. Fixing a basepoint $z_0 \in \mathcal{S}$, we can choose R and a uniformization $G : \mathbb{D}_R \to \mathcal{S}$ satisfying $G(0) = z_0, (\pi \circ G)'(0) = 1$, and it is easy to see that R and G are then uniquely determined.

Definition 2.5. *For $(\mathcal{S}, z_0)$ a pointed log-Riemann surface such that $\mathcal{S}^\times$ is simply connected, the conformal radius R of $(\mathcal{S}, z_0)$ is the unique $0 < R \leq \infty$ as determined above.*

2.2. **Examples.**

We briefly describe some examples of log-Riemann surfaces:

0. Log-Riemann surfaces associated to the complex plane: Let $\mathcal{S} = \mathbb{C}$ and $\pi : \mathcal{S} \to \mathbb{C}$ be any automorphism of the plane $\pi(z) = az + b$, $a \in \mathbb{C}^*$, $b \in \mathbb{C}$. The associated metric space is the euclidean plane with the euclidean metric scaled by $|a|$. The ramification locus $\mathcal{R}$ is empty. It is easy to see that these are the only (connected) log-Riemann surfaces with $\mathcal{R} = \emptyset$.

1. The log-Riemann surface of the nth root: Let $\mathcal{S} = \mathbb{C} - \{0\}$ and $\pi : \mathcal{S} \to \mathbb{C}, \pi(z) = z^n$. It is easy to see that $(\mathcal{S}, \pi)$ is a log-Riemann surface isometric to the union of n slit planes $\mathbb{C} - [-\infty, 0]$ pasted isometrically along the $2n$ "sides" of the slits, with a single ramification point p of order n, $\overline{\mathcal{S}} = \mathcal{S}^\times = \mathcal{S} \cup \{p\}$, such that $\pi(p) = 0$.

2. Polynomial log-Riemann surfaces: Generalizing the previous example, let π be a polynomial with C the set of critical points. Then $\mathcal{S} = \mathbb{C} - C$ is a log-Riemann surface with $\#C$-many ramification points whose orders add up to the degree of π.

3. The log-Riemann surface of the logarithm: Let $\mathcal{S} = \mathbb{C} - \{0\}$ and $\pi : \mathcal{S} \to \mathbb{C}, \pi(z) = e^z$. It is easy to see that $(\mathcal{S}, \pi)$ is a log-Riemann surface isometric to the union of infinitely many slit planes $\mathbb{C} - [-\infty, 0]$ pasted isometrically along the slits, with a single ramification point p, of infinite order, $\overline{\mathcal{S}} = \mathcal{S} \cup \{p\}$, such that $\pi(p) = 0$.

4. The universal covering of a log-Riemann surface: Let $\pi : \tilde{\mathcal{S}} \to \mathcal{S}$ be the universal covering of a log-Riemann surface $\mathcal{S}$ with projection mapping $\pi_{\mathcal{S}} : \mathcal{S} \to \mathbb{C}$. The map $\pi_{\tilde{\mathcal{S}}} := \pi_{\mathcal{S}} \circ \pi$ is a local holomorphic diffeomorphism. With respect to the induced path metric on $\tilde{\mathcal{S}}$ the map π is a 1-Lipschitz local isometry and hence extends to a map between the completions $\pi : \tilde{\mathcal{S}}^* \to \mathcal{S}^*$. Moreover since $\mathcal{R}$ is discrete, $\tilde{\mathcal{S}}^* - \tilde{\mathcal{S}} = \pi^{-1}(\mathcal{R})$ is discrete, and hence endows $\tilde{\mathcal{S}}$ with a log-Riemann surface structure.

3. Caratheodory Theorem for log-Riemann surfaces

3.1. **Caratheodory convergence of log-Riemann surfaces.**

Recall that log-Riemann surfaces are endowed with the flat metric $|d\pi|$.

Definition 3.1. *A pointed sequence of log-Riemann surfaces* $(\mathcal{S}_n, z_n)$ *converges to a pointed log Riemann surface* $(\mathcal{S}, z_0)$ *if for any compact* $K \subset \mathcal{S}$ *containing* z_0 *there exists* $N = N(K) \geq 1$ *such that for* $n \geq N$ *there is an isometric embedding* ι *of* K *into* $\mathcal{S}_n$ *mapping* z_0 *into* z_n *such that* ι *is a translation in the charts* π, π_n *(the translation that maps* $\pi(z_0)$ *to* $\pi(z_n)$*).*

Example. Let $(\mathcal{S}, z_0)$ be a pointed log-Riemann surface with projection mapping $\pi : \mathcal{S} \to \mathbb{C}$, $\pi(z_0) = 0$. For $n \geq 1$ we consider the new log-Riemann surface $\mathcal{S}_n$ with projection mapping $\pi_n = n \cdot \pi$ and $z_n = z_0$. When $n \to +\infty$, we have that $(\mathcal{S}_n, z_n)$ has the planar log-Riemann surface as Caratheodory limit.

The Caratheodory limit of a sequence $(\mathcal{S}_n, z_n)$ if it exists is unique up to isometry:

Proposition 3.2. *Let* $(\mathcal{S}, z_0)$ *and* $(\mathcal{S}', z_0')$ *be two pointed log-Riemann surfaces both of which are Caratheodory limits of a sequence of pointed log-Riemann surfaces* $(\mathcal{S}_n, z_n)$*. Then there is an isometry* $T : \mathcal{S} \to \mathcal{S}'$ *taking* z_0 *to* z_0' *whose expression in log-charts is the translation mapping* $\pi(z_0)$ *to* $\pi'(z_0')$*.*

Proof: Consider the germ of holomorphic diffeomorphism T from $\mathcal{S}$ to $\mathcal{S}'$ mapping z_0 to z_0' whose expression in log-charts is the translation maping $\pi(z_0)$ to $\pi'(z_0')$.

Let $\gamma : [a, b] \to \mathcal{S}$ be a curve in $\mathcal{S}$ starting from z_0 along which T can be continued. For n large enough, if ι and ι' denote the isometric embeddings of γ and $T(\gamma)$ respectively into $\mathcal{S}_n$, on γ we must have

$$\iota = \iota' \circ T$$

Lemma 3.3. *The germ* T *can be continued analytically along all paths in* $\mathcal{S}$*.*

Proof: Suppose there is a path $\gamma : [a, b] \to \mathcal{S}$, $\gamma(a) = z_0, \gamma(b) = z \in \mathcal{S}$ such that T can be continued analytically along $\gamma([a, b))$ but not up to $\gamma(b) = z$. Since T is a local isometry, the limit $z' = \lim_{x \to b} T(\gamma(x))$ exists and must be a ramification point of order $n > 1$ since T cannot be continued to $\gamma(b)$.

Take $\delta > 0$ small enough so that $B(z', \delta)$ contains no other ramification points and so that $\overline{B(z, \delta)} \subset \mathcal{S}$. Let $b_1 \in [a, b)$ be such that $T(\gamma(b_1)) \in B(z', \delta)$. Let $\alpha : [c, d] \to \mathcal{S}$ be a circular loop in $\mathcal{S}$ that winds once around z, starting from $\gamma(b_1)$, $\alpha(c) = \gamma(b_1)$. We note that T can be continued along α, but $T(\alpha(c))$ is not equal to $T(\alpha(d))$.

For n large enough the compacts $\gamma([a, b]) \cup \alpha([c, d]) \cup \overline{B(z, \delta)} \subset \mathcal{S}$ and $T(\gamma([a, b_1])) \cup T(\alpha([c, d])) \subset \mathcal{S}'$ both embed isometrically into $\mathcal{S}_n$, let ι, ι' be the respective embeddings. Now, the ball $\iota(B(z, \delta))$ is completely contained in $\mathcal{S}_n$, so for the curve α we have

$$\iota(\alpha(c)) = \iota(\alpha(d)) = \iota(\gamma(b_1)) \quad \text{where } \alpha(c) = \alpha(d) = \gamma(b_1),$$

which implies

$$\iota'(T(\alpha(c))) = \iota'(T(\alpha(d)))$$

and hence, since ι' is an isometry,

$$T(\alpha(c)) = T(\alpha(d)),$$

a contradiction. $\diamond$

Lemma 3.4. *The continuation of T to all of $\mathcal{S}$ is single-valued.*

Proof: Let $\gamma : [a,b] \to \mathcal{S}$, $\gamma(a) = \gamma(b) = z_0$, be a closed path in $\mathcal{S}$. Consider the curve $T(\gamma) \in \mathcal{S}'$ given by continuing T along γ. Take n large enough so that the compacts $\gamma([a,b]) \subset \mathcal{S}$ and $T(\gamma([a,b])) \subset \mathcal{S}'$ both embed isometrically into $\mathcal{S}_n$ via isometries ι and ι' respectively. As before, along γ we have

$$\iota = \iota' \circ T$$

Since

$$\iota(\gamma(a)) = \iota(\gamma(b)) = \iota(z_0) = z_n,$$

it follows that

$$\iota'(T(\gamma(a))) = \iota'(T(\gamma(b)))$$

and hence

$$T(\gamma(a))) = T(\gamma(b))),$$

$\diamond$

It follows from the above lemmas that we obtain a globally defined map $T : \mathcal{S} \to \mathcal{S}'$. Applying the same arguments to the germ $S = T^{-1}$ given by the inverse of the initial germ T gives a map $S : \mathcal{S}' \to \mathcal{S}$, and it is straightforward to check that T and S define global mutual inverses. The conclusion of the Proposition follows. $\diamond$

3.2. **Convergence of uniformizations. Proof of Theorem 1.1 :** Let $C \subset \mathbb{D}_R$ be the image under F of the finite ramification points of $\mathcal{S}$. Then C is discrete and the complement $\mathbb{D}_R - C$ contains a ball around the origin. For any compact $K \subset \mathcal{S}$ containing z_0, for n large enough let $\iota_n : K \to \mathcal{S}_n$ be the corresponding embeddings. The maps $g_n = F_n \circ \iota_n \circ F^{-1}$ are well-defined and univalent on any compact in $\mathbb{D}_R - C$ for n large enough, satisfying $g_n(0) = 0, g_n'(0) = 1$. It follows from classical univalent function theory that they form a normal family on any simply connected subdomain of $\mathbb{D}_R - C$ containing the origin. Since normality is a local property and every point in $\mathbb{D}_R - C$ has a neighbourhood contained in such a simply connected subdomain, it follows that the family (g_n) is normal on $\mathbb{D}_R - C$. Let g be a normal limit on $\mathbb{D}_R - C$ of the g_n's; then g is univalent and $g(0) = 0$. Hence for any Jordan curve $\gamma \subset \mathbb{D}_R - C$ if 0 is in the bounded component of $\mathbb{C} - \gamma$ then 0 is in the bounded component of $\mathbb{C} - g(\gamma)$. It follows that g is bounded on the bounded component of $\mathbb{C} - \gamma$, and hence the singularities of g at the points of C are removable.

It is easy to see that the extension of g to $\mathbb{D}_R$ is also univalent; since $g(0) = 0, g'(0) = 1$, if $R = \infty$ then g must be the identity. If $R < \infty$, since $\limsup R_n \leq R$, g takes values in $\overline{\mathbb{D}_R}$, so by the Schwarz lemma g is the identity. So the only limit point of the sequence g_n is the identity, and the sequence converges to the identity. Hence F_n converges to F uniformly on compacts, in the sense that $F_n \circ \iota_n = g_n \circ F$

converges on compacts of $\mathcal{S}$ to F. If $\liminf R_n < R$, then along a subsequence the maps g_n take values in a disc compactly contained in $\mathbb{D}_R$, contradicting the convergence of g_n to the identity; hence $\liminf R_n \geq R$, so $\lim R_n = R$. $\diamond$

Proof of Theorem 1.2: In the notation as above, $G_n = F_n^{-1}, G = F^{-1}$, and we have seen that the univalent maps $g_n = F_n \circ \iota_n \circ F^{-1} \to id$ uniformly on compacts in $\mathbb{D}_R - C$, $\lim R_n = R$. It follows that $g_n^{-1} \to id$ and so $(F_n \circ \iota_n)^{-1} = G \circ g_n^{-1} \to G$ uniformly on compacts in $\mathbb{D}_R - C$. Also for any compact in $\mathbb{D}_R - C$ for n large we can write

$$\pi_n \circ G_n = (\pi_n \circ \iota_n) \circ (F_n \circ \iota_n)^{-1}$$

By the hypothesis on the isometric embeddings, $\pi_n \circ \iota_n = T_n \circ \pi$ where T_n is the translation sending $\pi(z_0)$ to $\pi_n(z_n)$; since $\pi_n(z_n) \to \pi(z_0)$, $T_n \to id$, so $\pi_n \circ \iota_n \to \pi$ uniformly on compacts in $\mathcal{S}$, hence $\pi_n \circ G_n \to \pi \circ G$ uniformly on compacts in $\mathbb{D}_R - C$, and hence by the maximum principle also on $\mathbb{D}_R$. $\diamond$

Proof of Corollary 1.3: Let $\mathcal{S}_n$ be the log-Riemann surface of the nth root with projection $\pi_n : \mathcal{S}_n \to \mathbb{C}$, and take basepoints z_n such that $\pi_n(z_n) = 1$. Let $\mathcal{S}$ be the log-Rieman surface of the logarithm with projection $\pi : \mathcal{S} \to \mathbb{C}$ and basepoint z_0 such that $\pi(z_0) = 1$. Then any compact K in $\mathcal{S}$ can only intersect finitely many "sheets" of $\mathcal{S}$ and hence embeds isometrically in $\mathcal{S}_n$ for n large enough, so $(\mathcal{S}_n, z_n) \to (\mathcal{S}, z_0)$. The normalized uniformizations G_n, G of $\mathcal{S}_n, \mathcal{S}$ are given in charts by $\pi_n \circ G_n(z) = (1 + \frac{z}{n})^n$ and $\pi \circ G(z) = e^z$ respectively so the previous Theorem gives the conclusion of the Corollary. $\diamond$

References

[Bis05] K. Biswas. Smooth combs inside hedgehogs. *DISCRETE AND CONTINUOUS DYNAMICAL SYSTEMS, vol. 12, 5*, pages 853–880, 2005.

[Bis08] K. Biswas. Hedgehogs of hausdorff dimension one. *Ergodic Theory and Dynamical Systems, vol. 28, 6*, pages 1713–1727, 2008.

[BPM10] K. Biswas and R. Perez-Marco. Uniformization of simply connected finite type log-riemann surfaces. *preprint*, 2010.

[PM93] R. Perez-Marco. Sur les dynamiques holomorphes non-linearisables et une conjecture de v.i. arnold. *Annales Scientifiques de l'E.N.S 26*, pages 565–644, 1993.

[PM95] R. Perez-Marco. Uncountable number of symmetries for non-linearisable holomorphic dynamics. *Inventiones Mathematicae 119*, pages 67–127, 1995.

[PM00] R. Perez-Marco. Siegel disks with smooth boundary. *preprint*, 2000.

RKM Vivekananda University, Belur Math, WB-711 202, India

CNRS, LAGA, UMR 7539, Université Paris 13, Villetaneuse, France